



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

associated before they left for the South. Mr. W. H. Hoyt tells me that three Starlings have been taken recently from a flock that is spending the winter on Noroton Hill, near Stamford. — LOUIS B. BISHOP, M. D., *New Haven, Conn.*

Snow Bunting at Sea.—About noon on November 17, 1900, a Snow Bunting (*Plectrophenax nivalis*) was observed at sea by Mr. Paul du Chaillu and myself, on board the American line steamer 'New York.' The bird approached from a southwesterly direction and alighted in the rigging, flying in a rather labored manner. The vessel was then 260 miles east of Sandy Hook and 50 miles distant from the eastern end of Nantucket, the nearest land. The weather was fair. — HUGH M. SMITH, M. D., *Washington, D. C.*

Montana Redpolls.—A small series of Redpolls collected by Mr. Charles T. Hodges at Miles City and Fort Keogh, Montana, during the winter of 1899-1900 has come recently into my possession. The majority are specimens of *Acanthis linaria*, but there are three skins of *A. l. rostrata*, two of *A. l. holbællii*, and two of *A. hornemannii exilipes*.

The Greater Redpoll was taken on March 1 and 6, the Holbæll's (which Prof. Ridgway has examined) on March 2, and the Hoary, on February 26 and March 12. I can find previous records for only *Acanthis linaria* from Montana. — LOUIS B. BISHOP, M. D., *New Haven, Conn.*

***Acanthis linaria rostrata* and *Xanthocephalus xanthocephalus* in Connecticut.**—In a small collection of bird skins given me some time ago by Dr. W. H. Hotchkiss of this city I find two skins of the Greater Redpoll. They were taken by Dr. Hotchkiss near New Haven on December 17, 1878, and are, I believe, the only specimens of this subspecies so far recorded from Connecticut.

In the same collection was an unlabelled skin of a female Yellow-headed Blackbird, which Dr. Hotchkiss told me he was certain was shot near New Haven in June, 1878. In this connection it may be well to report that another female of this species was taken on Monomoy Island, Mass., September 8, 1897, by Mr. W. B. Revere, and given to me while in the flesh. — LOUIS B. BISHOP, M. D., *New Haven, Conn.*

Deformity of Maxilla in the House Sparrow.—The accompanying illustrations show the overgrowth of maxilla after loss of mandible in a male *Passer domesticus*. The photograph was taken by Mr. A. H. Verrill, who shot the bird in his yard in New Haven, December 10, 1900, and brought it to me in the flesh.

The culmen measures .60 inch from nostril against an average of .39 inch in five normal males of this species. The maxilla measures .16 inch in depth at tip on the right side, but had been worn off on the left where all that was left of the mandible approached it. The normal outer cov-